

September 2009

Bubbling Over: Soda Consumption and Its Link to Obesity in California

Susan H. Babey, Malia Jones, Hongjian Yu and Harold Goldstein

In California, 62% of adolescents ages 12-17 and 41% of children ages 2-11 drink at least one soda or other sweetened beverage every day. In addition, 24% of adults drink at least one soda or other sweetened beverage on an average day. Adults who drink soda occasionally (not every day) are 15% more likely to be overweight or obese, and adults who drink one or more sodas per day are 27% more likely to be overweight or obese than adults who do not drink soda, even when adjusting for poverty status and race/ethnicity.

This policy brief, produced collaboratively by the California Center for Public Health Advocacy and the UCLA Center for Health Policy Research, examines soda consumption in California by cities and counties using data from the 2005 California Health Interview Survey (CHIS 2005). In addition, the brief investigates whether there is an association between soda consumption and the prevalence of overweight and obesity.

There are major differences in soda consumption rates by geographic area in California, suggesting that social and environmental factors affect the consumption of soda. Also, the prevalence of overweight and obesity is higher among those who drink one or more sodas or other sweetened beverages every day than among those who do not consume these soft drinks. Establishing public policies that focus on reducing soda consumption could contribute to reversing California's increasing overweight and obesity problem.

Background

The prevalence of overweight and obesity has increased dramatically in both adults and children in the last three decades in the

United States. In the 1970s, about 15% of adults were obese and by 2004 the rate had climbed to 32%.¹ Although the prevalence of overweight among children is lower than among adults, the rates among children and adolescents have increased considerably more. The prevalence of overweight and obesity nearly tripled among 12-19 year olds and more than quadrupled among 6-11 year olds in the last three decades.

In California, 21% of adults are currently obese and an additional 35% are overweight. Among adolescents, 14% are obese and another 16% are overweight.² Similar to national trends, the trend in California is toward increasing weight in both adults and adolescents.³ Each year in California, overweight and obesity cost families, employers, the health care industry and the government \$21 billion.⁴ California spends more public and private money on the health consequences of obesity than any other state.⁵

Overweight and obesity are associated with serious health risks. In children and adolescents, overweight and obesity are associated with increased risk for cardiovascular disease indicators including



This policy brief was developed in collaboration with the California Center for Public Health Advocacy

high total cholesterol, high blood pressure, and high fasting insulin, an early indicator of diabetes risk.⁶ In addition, overweight children and adolescents are more likely to be overweight or obese as adults.⁷ In adults, overweight and obesity are associated with increased risk for diabetes, heart disease, stroke, some types of cancer and premature death.^{1, 8, 9}

Drinking sweetened beverages such as soda and fruit drinks that have added caloric sweeteners (e.g., sucrose, high fructose corn syrup) is one marker of a poor diet, and is associated with overweight and obesity in people of all ages.¹⁰⁻¹³ A number of studies have found that greater consumption of sweetened beverages is associated with overweight and obesity among both adults and children.¹²⁻¹⁹ In addition, randomized controlled trials that examine the impact of reducing intake of sweetened beverages on weight indicate that reducing consumption of soda and other sweetened drinks leads to reductions in overweight and obesity.^{20, 21} Among adults, drinking soda is also associated with increased risk for type 2 diabetes.¹³

Moreover, drinking sweetened beverages has increased, and it is now more common than ever, particularly among adolescents.²² Between 1977 and 2002 Americans increased their calorie intake from soft drinks by 228%.²³ Portion sizes have also increased from an average serving size of 6.5 fl oz (88 calories) in the 1950s, to 12 fl oz (150 calories), 20 fl oz (266 calories), and even larger portion sizes common today.²⁴⁻²⁶ The average serving size of soft drinks in fast food restaurants in 2002 was 23 fl oz (299 calories), with some chains now commonly selling soft drinks in 32 to 64 fl oz portions (416 to 832 calories, respectively).²⁷ Sweetened beverages are a significant contributor to total caloric intake, especially for children and adolescents, and they lack the nutrients our bodies need.^{24, 26, 28} Additionally, eating habits established in childhood are important determinants of eating habits as adults.^{29, 30}

Soda Consumption in California

Drinking sweetened beverages is common among California adults, adolescents and children. Data from CHIS 2005 show that nearly one out of four adults (24%) drink at least one soda every day—6.4 million California adults—and 36% drink soda occasionally, but not every day. Forty percent of adults report not drinking soda at all. In addition, 41% of children ages 2-11 drink at least one soda every day, nearly 2.2 million children in all. The rates of soda consumption among adolescents are much higher than among adults or children. More than 62% of adolescents ages 12-17—over two million teens—drink soda every day, including 13% (over 400,000) who drink three or more sodas every day. California adolescents drink 1.2 sodas per day on average. Conservatively assuming one soda is a 12-ounce can which contains 10 teaspoons of sugar, the average California adolescent consumes the equivalent of 39 pounds of sugar each year from soda and other sweetened beverages.

Soda Consumption Associated with Higher Prevalence of Overweight and Obesity

In California, 56% of adults and 30% of adolescents are either overweight or obese. The prevalence of overweight and obesity is higher among adults and adolescents who drink soda than among those who don't.

For both adults and adolescents, rates of overweight and obesity are 18% higher among those who drink one or more sodas every day compared to those who do not drink soda. Among adults, 62% of those who drink one or more sodas daily are either overweight or obese compared to 52% of adults who do not drink soda. Among adolescents, 32% of those who consume at least one soda per day are either overweight or obese, while 27% of those who consume no sodas on a typical day are either overweight or obese.

Soda consumption is associated with poverty and race/ethnicity; lower income people and people of color tend to drink more soda.³¹ These same groups also tend to be at higher risk for overweight and obesity. However, in our analysis of California adults, the association between soda consumption and overweight or obesity was independent of poverty status and race/ethnicity. Adults who drink soda occasionally (not every day) are 15% more likely to be overweight or obese, and adults who drink one or more sodas per day are 27% more likely to be overweight or obese than adults who do not drink soda, even when adjusting for poverty status and race/ethnicity (Exhibit 1).


Among adolescents, the association between soda consumption and overweight is not independent of poverty status and race/ethnicity. This may be partially due to the relatively small sample size for adolescents compared to adults. Compared to white adolescents, African-American and Latino adolescents are more likely to consume soda daily, while Asian adolescents are less likely. Adolescents from lower-income families are more likely to drink soda every day compared to adolescents from higher-income families.

Soda Consumption Varies from Place to Place in California

Trends in soda consumption and obesity may be influenced by social and environmental factors. For example, the food environment, including the presence of fast-food outlets, convenience stores, grocery stores and other food vendors, has an impact on health and dietary choices of the local population.³²⁻³⁴ A recent study by the California Center for Public Health Advocacy showed that California has more than four times as many fast-food restaurants and convenience stores as grocery stores and produce vendors—suggesting that Californians have greater access to foods with lower nutritional values than to healthier foods.³⁵ Moreover, this food environment has been linked to the prevalence of obesity and diabetes among California adults.³²

Increased Likelihood of Being Overweight or Obese for Those Who Drink Sodas Compared to Those Who Do Not, Adjusted for Race/Ethnicity and Income, Adults Age 18 and Over, California, 2005

Exhibit 1



Source: 2005 California Health Interview Survey

At the same time, soda consumption is associated with the use of fast-food restaurants among adolescents, and there is wide variation in the relative availability of fast-food restaurants in California communities.^{32, 36, 37}

Findings from CHIS 2005 show that there are major geographic differences in soda consumption in California (Exhibit 2). The percent of children drinking at least one soda each day ranges from 18% in Marin County to 61% in Imperial County. Among adolescents, the percent drinking one or more sodas each day ranges from 39% in Mendocino County to 78% in San Joaquin County. Among adults, the percent drinking one or more sodas each day ranges from just 11% in Marin County to 39% in Kings County.

Soda consumption also varies considerably among cities and census designated places (Exhibit 3). Among children and adolescents ages 2-17, the percent drinking at least one

Exhibit 2

Percent Drinking One or More Sodas per Day by County or County Group, Children, Adolescents and Adults, California, 2005

	Children Ages 2–11	Adolescents Ages 12–17	Adults Age 18 and Over
	One or More Sodas %	One or More Sodas %	One or More Sodas %
California	41.2	62.2	24.3
Alameda	31.0	58.9	17.4
Butte	30.4	61.8	20.3
Contra Costa	40.7	47.2	21.2
Del Norte, Siskiyou, Lassen, Trinity, Modoc, Plumas, Sierra	24.5	63.0	20.8
El Dorado	31.8	55.3	21.6
Fresno	53.1	68.7	35.0
Humboldt	33.2	50.3	16.4
Imperial	60.7	61.2	36.4
Kern	55.0	67.2	36.6
Kings	57.2	57.7	39.1
Lake	31.6	62.8	30.1
Los Angeles	44.3	64.9	25.5
Madera	39.9	75.3	37.4
Marin	18.4	41.3	10.6
Mendocino	38.1	39.0	18.8
Merced	55.4	*	32.7
Monterey	32.8	58.1	27.1
Napa	41.5	56.8	27.3
Nevada	25.6	40.9	17.5
Orange	36.9	56.4	23.4
Placer	31.5	66.2	18.4
Riverside	40.6	69.5	29.5
Sacramento	35.4	55.5	23.6
San Benito	26.4	58.9	25.6
San Bernardino	49.6	68.5	29.6
San Diego	34.8	63.1	21.1
San Francisco	21.5	42.1	10.9
San Joaquin	44.2	77.8	26.6
San Luis Obispo	41.7	66.8	18.3
San Mateo	32.5	50.1	14.4
Santa Barbara	39.8	53.8	19.0
Santa Clara	40.9	48.2	21.1
Santa Cruz	41.4	56.0	15.5
Shasta	32.0	60.0	27.5
Solano	45.2	58.7	26.1
Sonoma	42.0	60.7	20.7
Stanislaus	47.5	*	34.3
Sutter	44.5	*	29.2
Tehama, Glenn, Colusa	36.8	*	30.1
Tuolumne, Calaveras, Amador, Inyo, Mariposa, Mono, Alpine	35.0	*	17.3
Tulare	44.2	71.0	36.1
Ventura	39.0	60.4	24.8
Yolo	37.3	62.4	13.9
Yuba	50.5	62.9	30.9

Note:

* Indicates the estimate was not statistically reliable. Not all differences between rates are statistically significant. The 95% confidence intervals are available at: http://www.healthpolicy.ucla.edu/soda_consumption.html

Source: 2005 California Health Interview Survey

Percent Drinking One or More Sodas per Day by Cities and Census Designated Places,
California, 2005

Exhibit 3

	Children and Adolescents Ages 2–17 %	Adults Age 18 and Over %		Children and Adolescents Ages 2–17 %	Adults Age 18 and Over %
California	49.4	24.3	Mission Viejo	43.3	18.0
Anaheim	45.4	26.5	Modesto	57.0	31.8
Antioch	44.8	21.9	Moreno Valley	55.4	33.7
Bakersfield	60.1	33.9	Murrieta	49.7	26.5
Baldwin Park	52.2	29.0	Norwalk	51.5	31.0
Bellflower	51.3	30.9	Oakland	44.1	20.6
Buena Park	44.0	24.5	Oceanside	47.7	20.8
Burbank	48.3	19.6	Ontario	57.7	32.9
Carlsbad	43.5	16.3	Orange	46.0	22.6
Carson	52.7	25.0	Oxnard	50.6	30.0
Chino	56.3	31.2	Palmdale	54.9	32.1
Chino Hills	52.4	22.2	Pasadena	54.2	22.9
Chula Vista	46.2	23.1	Pomona	56.6	29.5
Citrus Heights	39.4	21.9	Rancho Cucamonga	54.6	26.0
Clovis	53.8	27.0	Redding	44.2	25.3
Compton	54.7	33.2	Rialto	59.4	32.8
Concord	44.2	21.5	Richmond	46.1	28.4
Corona	50.7	29.6	Riverside	49.8	31.7
Costa Mesa	43.5	25.0	Roseville	43.6	16.4
Daly City	38.3	13.7	Sacramento	44.3	25.4
Downey	51.4	29.6	Salinas	46.9	28.9
East Los Angeles *	53.3	38.4	San Bernardino	58.6	32.7
El Cajon	47.6	22.2	San Buenaventura (Ventura)	46.6	22.3
El Monte	51.8	29.2	San Diego	46.2	22.8
Elk Grove *	43.3	21.2	San Francisco	36.9	11.5
Escondido	48.1	22.6	San Jose	42.8	21.7
Fairfield	47.0	26.5	Santa Ana	47.3	33.2
Florence-Graham *	54.2	36.5	Santa Clara	40.6	19.2
Fontana	57.5	31.9	Santa Clarita	49.9	20.6
Fremont	38.0	14.1	Santa Maria	48.3	24.1
Fresno	57.4	33.5	Santa Rosa	45.4	19.7
Fullerton	44.0	23.6	Simi Valley	44.0	20.5
Garden Grove	43.9	24.0	Southgate	52.9	36.8
Glendale	47.6	19.6	Stockton	57.3	28.1
Hawthorne	53.2	31.4	Sunnyvale	39.8	18.7
Hayward	41.3	18.4	Temecula	47.8	28.2
Hesperia	55.5	27.2	Thousand Oaks	43.8	19.8
Huntington Beach	40.7	20.7	Torrance	46.0	18.9
Indio	55.6	37.5	Tracy	56.9	24.9
Inglewood	55.0	32.6	Vacaville	45.4	25.4
Irvine	43.6	19.5	Vallejo	48.8	25.7
Lancaster	54.8	30.7	Victorville	57.0	29.2
Livermore	41.1	15.1	Visalia	56.3	30.8
Long Beach	51.5	27.2	Vista	48.8	23.8
Los Angeles	51.9	24.8	West Covina	50.4	21.6
Lynwood	53.5	33.3	Westminster	42.8	22.4
Merced	61.9	33.3			

Note:

* Indicates a Census Designated Place. Census designated places are communities that lack separate governments but otherwise resemble incorporated places such as cities. This table includes only cities in which the combined population of children and adolescents ages 2-17 was at least 20,000. Not all differences between rates are statistically significant. The 95% confidence intervals are available at: http://www.healthpolicy.ucla.edu/soda_consumption.html

Source: 2005 California Health Interview Survey

soda per day ranged from 37% in San Francisco to 62% in Merced. Among adults, the percent drinking at least one soda per day ranged from 12% in San Francisco to 38% in East Los Angeles.

Conclusions

In California, 62% of adolescents ages 12-17 and 41% of children ages 2-11 drink at least one soda or other sweetened beverage every day. In addition, nearly one out of four adults (24%) drink soda every day and 36% drink soda occasionally. This amounts to 10.7 million Californians over the age of one who drink at least one soda each day. This soda consumption greatly increases the amount of added sugar and other caloric sweeteners in the diet of Californians without contributing substantially to the nutritional needs of the population.

For both adults and adolescents, the prevalence of overweight and obesity is higher among those who drink one or more sodas or other sweetened beverages every day than among those who do not. Among adults, even after adjusting for race and household income, those who drink one or more sodas each day are 27% more likely to be overweight or obese than adults who do not drink soda. These findings are consistent with other research.³⁸ Additionally, childhood eating habits and weight status are important determinants of health as adults.^{7, 29, 30} Taken together, these findings suggest a number of potential benefits from reducing soft drink consumption including reduced risk of obesity, improved dietary intake and reduced risk of diabetes.

Data Source and Methods

This policy brief examines geographical variation in soda consumption among children, adolescents and adults in California as well as its association with overweight and obesity among adults and adolescents using data from the 2005 California Health Interview Survey (CHIS 2005). All statements in this report that compare rates for one group with another group reflect statistically significant differences ($p < 0.05$) unless otherwise noted. CHIS 2005 completed interviews with over 4,000 adolescents and over 43,000 adults, drawn from every county in the state, in English, Spanish, Chinese (both Mandarin and Cantonese),

Vietnamese and Korean. The California Health Interview Survey is a collaboration of the UCLA Center for Health Policy Research, California Department of Public Health, the California Department of Health Care Services and the Public Health Institute. Funding for the CHIS 2005 statewide survey was provided by the California Department of Health Care Services, The California Endowment, the National Cancer Institute, The Robert Wood Johnson Foundation, the California Children and Families Commission, the California Office of the Patient Advocate, the California Department of Mental Health, the Centers for Disease Control and Prevention (CDC) and Kaiser Permanente. For local funders and other information on CHIS, visit www.chis.ucla.edu.

In adults, overweight is defined as a Body Mass Index (BMI) between 25 and 30, while obesity is defined as BMI of 30 or greater. Among adolescents, overweight is defined as having a BMI between the 85th and 95th percentile on the Centers for Disease Control and Prevention's BMI-for-age growth charts, while obesity is defined as having a BMI above the 95th percentile.³⁹

Adults and adolescents self-reported their consumption of soda and other sweetened beverages. Adults were asked the following two questions: "During the past month, how many times (per day, per week or per month) did you drink soda such as Coke or 7-Up? Do not include diet soda." and "How many times did you drink fruit-flavored drinks such as lemonade or Sunny Delight? Do not include diet drinks." Responses to these questions were combined and converted to a common metric to estimate daily consumption of soda and other sweetened beverages. Adolescents were asked: "Yesterday, how many glasses or cans of soda such as Coke, or other sweetened drinks such as fruit punch or Sunny Delight did you drink? Do not count diet drinks." For children ages 2-11, the most knowledgeable parent or guardian responded to the following question: "Yesterday, how many glasses or cans of soda such as Coke or other sweetened drinks such as fruit punch or Sunny Delight did (he/she) drink? Do not count diet drinks." For all respondents, consumption of 100% fruit juice was reported in a previous question and is not included in our estimates of sweetened beverage consumption.

We used small-area estimation to generate model-based estimates of the proportion of adults and children who consume one or more sodas per day for each city.^{40, 41} Small-area estimation uses modeling to produce estimates for small geographic areas, such as cities, for which there is not sufficient sample to produce direct estimates. The models are based on

individual-level demographic and health outcome data from CHIS 2005 as well as demographic data at the census block group level from the Census and Claritas Inc. To maximize the reliability and validity of the estimates, we present only estimates for cities with a population of at least 20,000 for the age group being modeled. For more information about small-area estimation methodology, see: Yu H, Meng YY, Mendez-Luck CA, Jhavar M, Wallace SP. *Small-Area Estimation of Health Insurance Coverage for California Legislative Districts*.

Author Information

Susan H. Babey, PhD, is a senior research scientist at the UCLA Center for Health Policy Research. Malia Jones, MPH, is a graduate student researcher at the UCLA Center for Health Policy Research. Hongjian Yu, PhD, is the director for statistical support at the UCLA Center for Health Policy Research. Harold Goldstein, DrPH, is the Executive Director of the California Center for Public Health Advocacy.

Acknowledgements

The authors wish to thank Theresa Hastert, MPP, Jenny Chia, PhD, Lijie Di, MS, Gwen Driscoll and Celeste Maglan for their assistance. The authors would also like to thank the following individuals for their helpful comments: Stefan Harvey, Assistant Director, the California Center for Public Health Advocacy; Lisa Hershey, MPH, Chief, Coordinating Office for Obesity Prevention, California Department of Public Health; Leslie Mikkelsen, MPH, Managing Director, Prevention Institute; Gail Woodward-Lopez, MPH, RD, Associate Director, Center for Weight and Health, University of California, Berkeley.

Funding

Support for this policy brief was provided by a grant from The California Endowment to the California Center for Public Health Advocacy.

Suggested Citation

Babey SH, Jones M, Yu H, Goldstein H. *Bubbling Over: Soda Consumption and Its Link to Obesity in California*. UCLA Center for Health Policy Research and California Center for Public Health Advocacy, 2009.

Endnotes

- 1 Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA*. Apr 5 2006;295(13):1549-1555.
- 2 2005 California Health Interview Survey.
- 3 Babey SH, Grant D, Brown ER. *Adult Smoking Rate Declines, While Asthma, Diabetes and Obesity Rates Rise*. Los Angeles: UCLA Center for Health Policy Research; Nov 2006.

- 4 *The Economic Costs of Overweight, Obesity, and Physical Inactivity Among California Adults – 2006* (July 2009). The California Center for Public Health Advocacy.
- 5 Finkelstein EA, Fiebelkorn IC, Wang G. State-level estimates of annual medical expenditures attributable to obesity. *Obesity Research*. Jan 2004;12(1):18-24.
- 6 Freedman DS, Mei Z, Srinivasan SR, Berenson GS, Dietz WH. Cardiovascular risk factors and excess adiposity among overweight children and adolescents: the Bogalusa Heart Study. *Journal of Pediatrics*. Jan 2007;150(1):12-17 e12.
- 7 Guo SS, Wu W, Chumlea WC, Roche AF. Predicting overweight and obesity in adulthood from body mass index values in childhood and adolescence. *American Journal of Clinical Nutrition*. Sep 2002;76(3):653-658.
- 8 Must A, Spadano J, Coakley EH, Field AE, Colditz G, Dietz WH. The disease burden associated with overweight and obesity. *JAMA*. Oct 27 1999;282(16):1523-1529.
- 9 Ogden CL, Yanovski SZ, Carroll MD, Flegal KM. The epidemiology of obesity. *Gastroenterology*. May 2007;132(6):2087-2102.
- 10 French SA, Lin B-H, Guthrie JF. National trends in soft drink consumption among children and adolescents age 6 to 17 years: Prevalence, amounts, and sources, 1977/1978 to 1994/1998. *Journal of the American Dietetic Association*. 2003;103(10):1326-1331.
- 11 Harnack L, Stang J, Story M. Soft Drink Consumption Among U.S. Children and Adolescents: Nutritional Consequences. *Journal of the American Dietetic Association*. 1999;99(4):436-441.
- 12 Malik VS, Schulze MB, Hu FB. Intake of sugar-sweetened beverages and weight gain: a systematic review. *American Journal of Clinical Nutrition*. Aug 2006;84(2):274-288.
- 13 Schulze MB, Manson JE, Ludwig DS, et al. Sugar-sweetened beverages, weight gain, and incidence of type 2 diabetes in young and middle-aged women. *JAMA*. Aug 25 2004;292(8):927-934.
- 14 Ludwig DS, Peterson KE, Gortmaker SL. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *The Lancet*. 2001;357(9255):505-508.
- 15 Giammattei J, Blix G, Marshak HH, Wollitzer AO, Pettitt DJ. Television watching and soft drink consumption: associations with obesity in 11- to 13-year-old schoolchildren. *Archives of Pediatric and Adolescent Medicine*. Sep 2003;157(9):882-886.
- 16 Phillips SM, Bandini LG, Naumova EN, et al. Energy-dense snack food intake in adolescence: longitudinal relationship to weight and fatness. *Obesity Research*. Mar 2004;12(3):461-472.
- 17 LaRowe TL, Moeller SM, Adams AK. Beverage patterns, diet quality, and body mass index of U.S. preschool and school-aged children. *Journal of the American Dietetic Association*. Jul 2007;107(7):1124-1133.
- 18 Welsh JA, Cogswell ME, Rogers S, Rockett H, Mei Z, Grummer-Strawn LM. Overweight among low-income preschool children associated with the consumption of sweet drinks: Missouri, 1999-2002. *Pediatrics*. Feb 2005;115(2):e223-229.
- 19 Dubois L, Farmer A, Girard M, Peterson K. Regular sugar-sweetened beverage consumption between meals increases risk of overweight among preschool-aged children. *Journal of the American Dietetic Association*. Jun 2007;107(6):924-934; discussion 934-925.
- 20 Ebbeling CB, Feldman HA, Osganian SK, Chomitz VR, Ellenbogen SJ, Ludwig DS. Effects of decreasing sugar-sweetened beverage consumption on body weight in adolescents: a randomized, controlled pilot study. *Pediatrics*. Mar 2006;117(3):673-680.



The UCLA Center
for Health Policy Research
is affiliated with the
UCLA School of Public Health
and the UCLA School of Public Affairs.

The views expressed in this policy brief
are those of the authors and do not
necessarily represent the UCLA Center for
Health Policy Research, the Regents of the
University of California, or collaborating
organizations or funders.

PB2009-5

Copyright © 2009 by the Regents of the
University of California and the California
Center for Public Health Advocacy

Editor-in-Chief: E. Richard Brown, PhD

Phone: 310-794-0909
Fax: 310-794-2686
Email: chpr@ucla.edu
Web Site: www.healthpolicy.ucla.edu

- 21 James J, Thomas P, Cavan D, Kerr D. Preventing childhood obesity by reducing consumption of carbonated drinks: cluster randomised controlled trial. *British Medical Journal*. May 22 2004;328(7450):1237.
- 22 Wang YC, Bleich SN, Gortmaker SL. Increasing caloric contribution from sugar-sweetened beverages and 100% fruit juices among U.S. children and adolescents, 1988-2004. *Pediatrics*. Jun 2008;121(6):e1604-1614.
- 23 Duffey KJ, Popkin BM. Shifts in patterns and consumption of beverages between 1965 and 2002. *Obesity (Silver Spring)*. Nov 2007;15(11):2739-2747.
- 24 Jacobsen M. Liquid candy: How soft drinks are harming Americans' health. <http://www.cspinet.org/liquidcandy/>. Accessed December 22, 2008.
- 25 Nielsen SJ, Popkin BM. Patterns and trends in food portion sizes, 1977-1998. *JAMA*. Jan 22-29 2003;289(4):450-453.
- 26 Nielsen SJ, Popkin BM. Changes in beverage intake between 1977 and 2001. *American Journal of Preventive Medicine*. Oct 2004;27(3):205-210.
- 27 Young LR, Nestle M. Expanding portion sizes in the U.S. marketplace: implications for nutrition counseling. *Journal of the American Dietetic Association*. Feb 2003;103(2):231-234.
- 28 Block G. Foods contributing to energy intake in the U.S.: data from NHANES III and NHANES 1999-2000. *Journal of Food Composition and Analysis*. 2004;17(3-4):439-447.
- 29 Mikkila V, Rasanen L, Raitakari OT, Pietinen P, Viikari J. Consistent dietary patterns identified from childhood to adulthood: the cardiovascular risk in Young Finns Study. *British Journal of Nutrition*. Jun 2005;93(6):923-931.
- 30 Lake AA, Mathers JC, Rugg-Gunn AJ, Adamson AJ. Longitudinal change in food habits between adolescence (11-12 years) and adulthood (32-33 years): the ASH30 Study. *Journal of Public Health*. Mar 2006;28(1):10-16.
- 31 Rehm CD, Matte TD, Van Wye G, Young C, Frieden TR. Demographic and behavioral factors associated with daily sugar-sweetened soda consumption in New York City adults. *Journal of Urban Health*. May 2008;85(3):375-385.
- 32 *Designed for Disease: the Link Between Local Food Environments and Obesity and Diabetes*: California Center for Public Health Advocacy, PolicyLink, and the UCLA Center for Health Policy Research.; April 2008.
- 33 Morland K, Diez Roux A, Wing S. Supermarkets, other food stores, and obesity: the atherosclerosis risk in communities study. *American Journal of Preventive Medicine*. Apr 2006;30(4):333-339.
- 34 Morland K, Wing S, Diez Roux A. The contextual effect of the local food environment on residents' diets: the atherosclerosis risk in communities study. *American Journal of Public Health*. Nov 2002;92(11):1761-1767.
- 35 *Searching for healthy food: The food landscape in California cities and counties*: California Center for Public Health Advocacy; 2007.
- 36 Taveras EM, Berkey CS, Rifas-Shiman SL, et al. Association of consumption of fried food away from home with body mass index and diet quality in older children and adolescents. *Pediatrics*. Oct 2005;116(4):e518-524.
- 37 Wiecha JL, Finkelstein D, Troped PJ, Fragala M, Peterson KE. School vending machine use and fast-food restaurant use are associated with sugar-sweetened beverage intake in youth. *Journal of the American Dietetic Association*. Oct 2006;106(10):1624-1630.
- 38 Vartanian LR, Schwartz MB, Brownell KD. Effects of Soft Drink Consumption on Nutrition and Health: A Systematic Review and Meta-Analysis. *American Journal of Public Health*. April 1, 2007 2007;97(4):667-675.
- 39 Centers for Disease Control and Prevention. Defining overweight and obesity. <http://www.cdc.gov/nccdphp/dnpa/obesity/defining.htm>. Accessed November 14, 2008.
- 40 Yu H, Meng YY, Mendez-Luck CA, Jhawar M, Wallace SP. Small-area estimation of health insurance coverage for California legislative districts. *American Journal of Public Health*. Apr 2007;97(4):731-737.
- 41 Mendez-Luck CA, Yu H, Meng YY, Jhawar M, Wallace SP. Estimating health conditions for small areas: asthma symptom prevalence for state legislative districts. *Health Services Research*. Dec 2007;42(6 Pt 2):2389-2409.